

12th January	Corbettmaths										
$4^0 + 4^{\frac{1}{2}} + 4^1 + 4^2$											
	Find x										
<table style="width: 100%; border-collapse: collapse;"> <tr><td style="padding: 2px;">Lowest</td><td style="padding: 2px;">68kg</td></tr> <tr><td style="padding: 2px;">Lower Quartile</td><td style="padding: 2px;">74kg</td></tr> <tr><td style="padding: 2px;">Median</td><td style="padding: 2px;">82kg</td></tr> <tr><td style="padding: 2px;">Upper Quartile</td><td style="padding: 2px;">88kg</td></tr> <tr><td style="padding: 2px;">Highest</td><td style="padding: 2px;">100kg</td></tr> </table>	Lowest	68kg	Lower Quartile	74kg	Median	82kg	Upper Quartile	88kg	Highest	100kg	
Lowest	68kg										
Lower Quartile	74kg										
Median	82kg										
Upper Quartile	88kg										
Highest	100kg										
<p>The students in a school sit two tests, a French test (F) and German test (G). Everyone passed at least one test. 78% passed the French test and 84% passed the German test.</p> <p>Show this information in the Venn diagram</p>	<p>£</p>										
<p>After a reduction of 6% in the original price, a caravan is sold for £7000.</p> <p>Both these values are correct to one significant figure.</p> <p>Calculate the lowest possible original price before the reduction was applied.</p>											

12th January	Corbettmaths										
$4^0 + 4^{\frac{1}{2}} + 4^1 + 4^2$											
	Find x										
<table style="width: 100%; border-collapse: collapse;"> <tr><td style="padding: 2px;">Lowest</td><td style="padding: 2px;">68kg</td></tr> <tr><td style="padding: 2px;">Lower Quartile</td><td style="padding: 2px;">74kg</td></tr> <tr><td style="padding: 2px;">Median</td><td style="padding: 2px;">82kg</td></tr> <tr><td style="padding: 2px;">Upper Quartile</td><td style="padding: 2px;">88kg</td></tr> <tr><td style="padding: 2px;">Highest</td><td style="padding: 2px;">100kg</td></tr> </table>	Lowest	68kg	Lower Quartile	74kg	Median	82kg	Upper Quartile	88kg	Highest	100kg	
Lowest	68kg										
Lower Quartile	74kg										
Median	82kg										
Upper Quartile	88kg										
Highest	100kg										
<p>The students in a school sit two tests, a French test (F) and German test (G). Everyone passed at least one test. 78% passed the French test and 84% passed the German test.</p> <p>Show this information in the Venn diagram</p>	<p>£</p>										
<p>After a reduction of 6% in the original price, a caravan is sold for £7000.</p> <p>Both these values are correct to one significant figure.</p> <p>Calculate the lowest possible original price before the reduction was applied.</p>											