

11th July		Corbettmaths
<p>Graph A</p>	<p>Graph B</p>	<p>$y = x^2$ is graph A</p> <p>$y = x^3$ is graph</p> <p>$y = 2^x$ is graph</p> <p>$y = \frac{1}{x}$ is graph</p>
<p>Graph C</p>	<p>Graph D</p>	
<p>For all the values of x</p> <p>$f(x) = x - 180$</p> <p>$g(x) = \cos x$</p> <p>Draw the graph of the function $y = gf(x)$ for $0^\circ \leq x \leq 360^\circ$</p>		
<p>A sequence of numbers is formed by the iterative process</p> <p>$a_{n+1} = (a_n)^2 - 10$</p> <p>$a_1 = 3$</p>	Find a_3	
<p>There are 20 sweets in a box. There are y chocolate sweets and the rest of the sweets are mints.</p> <p>Florence takes out two sweets, at random, from the box.</p>	Find an expression, in terms of y , for the probability that Florence takes two mint sweets.	

11th July		Corbettmaths
<p>Graph A</p>	<p>Graph B</p>	<p>$y = x^2$ is graph A</p> <p>$y = x^3$ is graph</p> <p>$y = 2^x$ is graph</p> <p>$y = \frac{1}{x}$ is graph</p>
<p>Graph C</p>	<p>Graph D</p>	
<p>For all the values of x</p> <p>$f(x) = x - 180$</p> <p>$g(x) = \cos x$</p> <p>Draw the graph of the function $y = gf(x)$ for $0^\circ \leq x \leq 360^\circ$</p>		
<p>A sequence of numbers is formed by the iterative process</p> <p>$a_{n+1} = (a_n)^2 - 10$</p> <p>$a_1 = 3$</p>	Find a_3	
<p>There are 20 sweets in a box. There are y chocolate sweets and the rest of the sweets are mints.</p> <p>Florence takes out two sweets, at random, from the box.</p>	Find an expression, in terms of y , for the probability that Florence takes two mint sweets.	